



PSICOLOGIA CLINICA

dott.ssa Paoletta Florio
psicologa-psicoterapeuta



DEFINIZIONE DI ALLEANZA TERAPEUTICA

alleanza terapeutica: il legame tra paziente e operatore e il reciproco accordo circa gli obiettivi del cambiamento e i compiti utili per realizzare tali obiettivi (Bordin, 1979)

- perché è utile?
- da chi dipende?
- come si favorisce?

A COSA SERVE?

- buon predittore degli esiti del trattamento
- favorisce l'aderenza al trattamento

- da chi dipende?
- come si favorisce?

DA CHI DIPENDE?

- paziente: si trova in condizione di bisogno e, talvolta, le caratteristiche stesse della sua patologia gli rendono difficile il compito di relazionarsi in modo adeguato e costruttivo
- operatore: si trova nella posizione di poter favorire la costruzione di una fruttuosa alleanza terapeutica e dovrebbe possedere le competenze necessarie
- **come si favorisce?**

COME SI FAVORISCE?



Pergamon

Clinical Psychology Review 23 (2003) 1–33

CLINICAL
PSYCHOLOGY
REVIEW

A review of therapist characteristics and techniques positively impacting the therapeutic alliance

Steven J. Ackerman^{a,*}, Mark J. Hilsenroth^b

^a*The Austen Riggs Center, P.O. Box 962, 25 Main Street, Stockbridge, MA 01262, USA*

^b*The Derner Institute of Advanced Psychological Studies, Adelphi University, Garden City, NY, USA*

Received 15 March 2001; received in revised form 26 July 2001; accepted 2 May 2002

Abstract

The present review is a comprehensive examination of the therapist's personal attributes and in-session activities that positively influence the therapeutic alliance from a broad range of psychotherapy perspectives. Therapist's personal attributes such as being flexible, honest, respectful, trustworthy, confident, warm, interested, and open were found to contribute positively to the alliance. Therapist techniques such as exploration, reflection, noting past therapy success, accurate interpretation, facilitating the expression of affect, and attending to the patient's experience were also found to contribute positively to the alliance. This review reveals how these therapist personal qualities and techniques have a positive influence on the identification or repair of ruptures in the alliance.

© 2003 Elsevier Science Ltd. All rights reserved.

COME SI FAVORISCE?

Journal of Psychiatric and Mental Health Nursing, 2012, 19, 1-7



The quality of the working alliance between chronic psychiatric patients and their case managers: process and outcomes

M. DE LEEUW¹ RN MSc CNS, B. VAN MEIJEL⁴ RN PhD,
M. GRYPDONCK^{2,5} RN PhD & H. KROON³ Mpsy PhD

¹Clinical Nurse Specialist in Mental Health, Altrecht Institute of Mental Healthcare, ²Professor of Nursing Science, The University of Utrecht, ³Director, The Trimbos Institute of Mental Health and Addiction, Utrecht, and

⁴Associate Professor of Mental Health Nursing, Research Group Mental Health Nursing, INHolland University for Applied Sciences, Amsterdam, the Netherlands, and ⁵Emeritus Professor of Nursing Science, The University of Ghent, Belgium

COME SI FAVORISCE?

De Leeuw et al. (2012)

caratteristiche della sintomatologia del paziente che inficiano la qualità dell'alleanza terapeutica:

- ostilità
- sfiducia
- alienazione
- abilità inadeguate di adattamento
- ridotta consapevolezza di malattia

COME SI FAVORISCE?

De Leeuw et al. (2012)

caratteristiche del case manager:

- fornisce supporto pratico nelle attività di vita quotidiana
- mostra empatia
- mostra rispetto
- attenzione sia alla «parte sana» che alla «parte malata»

COME SI FAVORISCE?

Pinto et al: Communication style and therapeutic alliance

Patient-centred communication is associated with positive
therapeutic alliance: a systematic review

Rafael Zambelli Pinto¹, Manuela L Ferreira¹, Vinicius C Oliveira², Marcia R Franco^{1,3},
Roger Adams², Christopher G Maher¹ and Paulo H Ferreira²

¹The George Institute for Global Health, University of Sydney, Australia, ²Discipline of Physiotherapy, Faculty of Health Sciences,
The University of Sydney, Australia, ³Regional Public Hospital of Betim, Minas Gerais, Brazil

Can J Psychiatry 2012;57(8):464-469

In Review

On the Self-Stigma of Mental Illness: Stages, Disclosure, and Strategies for Change

Patrick W Corrigan, PsyD¹; Deepa Rao, PhD, MA²

- public stigma: rappresenta il pregiudizio e la discriminazione diretta ad un gruppo e si riferisce agli atteggiamenti negativi nei confronti delle persone che vengono svalutate
- self-stigma: internalizzazione degli atteggiamenti pubblici

stereotipi negativi: caratterizzazioni (per esempio, pericolosità) delle persone che vivono certe condizioni da parte della società

pregiudizio: emozione (per esempio, paura) provata da chi condivide lo stereotipo negativo

discriminazione: comportamento che deriva dal pregiudizio; si manifesta in tre modi:

- perdita di opportunità (per esempio, non viene affittato un appartamento ad una persona con problemi di salute mentale)
- coercizione (per esempio, un'autorità prende decisioni per un paziente psichiatrico)
- segregazione (istituzionalizzazione o isolamento nei contesti sociali)

(Carozza, 2014)

il processo di auto-stigmatizzazione richiede:

- consapevolezza della discriminazione subita «chi ha un problema di salute mentale è pericoloso (o colpevole di essere malato e/o incompetente e/o parassita della società)»
- condivisione degli stereotipi da parte della persona malata «è vero»
- applicazione degli stereotipi a se stesso «sono pericoloso»


self-stigma:

«sono pericoloso»

«temo me stesso»

«mi isolo»

Figure 1 The stage model of self-stigma



impatto del self-stigma (Corrigan et al., 2009):

- riduce ridotta autostima e autoefficacia
- riduce la capacità di accedere ai servizi
- ostacola il raggiungimento degli obiettivi di vita
- determina esiti peggiori
- riduce la qualità di vita

Journal of Mental Health,
October 2008; 17(5): 482–491

informa
healthcare

Perceived stigma predicts low self-efficacy and poor coping in schizophrenia

BIRGIT KLEIM^{1,2}, ROLAND VAUTH^{1,3}, GUNDULA ADAM¹,
ROLF-DIETER STIEGLITZ³, PETER HAYWARD², &
PATRICK CORRIGAN⁴

¹Department of Psychiatry and Psychotherapy, University Hospital Freiburg, Germany, ²Institute of Psychiatry, Kings College London, UK, ³Psychiatric Outpatient Department, University Hospital Basel, Switzerland, and ⁴Illinois Institute of Technology, Chicago, IL, USA

Abstract

Background: The stigma of mental illness has been shown to be a barrier to recovery in schizophrenia.

Aims: The present study aimed to examine the influence of perceived stigma on self-efficacy and the coping styles of secrecy and withdrawal over and above selected sociodemographic and clinical variables and psychopathology.

Method: Outpatients diagnosed with schizophrenia ($N = 127$) filled in measures of perceived stigma, secrecy and withdrawal as coping modi, positive and negative symptom severity, self-efficacy, depression and insight into psychosis.

Results: Perceived stigma accounted for a significant amount of variance in secrecy and withdrawal, as well as low self-efficacy, over and above the amount of variance explained by positive and negative symptomatology, depression, insight, age and gender.

Conclusion: Our findings demonstrate the negative influence of stigma perceived by persons with schizophrenia on self-efficacy and coping. Clinicians should offer guidance in functional coping with perceived stigma in order to reduce the effect of stigma on patients' lives.

Connection between self-stigma, adherence to treatment, and discontinuation of medication

This article was published in the following Dove Press Journal:
Patient Preference and Adherence

22 July 2016

[Number of times this article has been viewed](#)

Dana Kamaradova¹
Klara Latalova¹
Jan Prasko¹
Radim Kubinek¹
Kristyna Vrbova¹
Barbora Mainerova¹
Andrea Cinculova¹
Marie Ociskova¹
Michaela Holubova²
Jarmila Smoldasova¹
Anezka Tichackova¹

¹Department of Psychiatry,
University Hospital Olomouc,

²Faculty of Medicine and Dentistry,
Palacky University Olomouc,
Olomouc, Czech Republic

Introduction: Self-stigma plays a role in many areas of the patient's life. Furthermore, it also discourages therapy. The aim of our study was to examine associations between self-stigma and adherence to treatment and discontinuation of medication in patients from various diagnostic groups.

Methods: This cross-sectional study involved outpatients attending the Department of Psychiatry, University Hospital Olomouc, Czech Republic. The level of self-stigma was measured with the Internalized Stigma of Mental Illness and adherence with the Drug Attitude Inventory. The patients also anonymously filled out a demographic questionnaire which included a question asking whether they had discontinued their medication in the past.

Results: We examined data from 332 patients from six basic diagnostic categories (substance abuse disorders, schizophrenia, bipolar disorders, depressive disorders, anxiety disorders, and personality disorders). The study showed a statistically significant negative correlation between self-stigma and adherence to treatment in all diagnostic groups. Self-stigma correlated positively and adherence negatively with the severity of disorders. Another important factor affecting both variables was partnership. Self-stigma positively correlated with doses of antidepressants and adherence with doses of anxiolytics. Self-stigma also negatively correlated with education, and positively with a number of hospitalizations and number of psychiatrists visited. Adherence was further positively correlated with age and age of onset of disorders. Regression analysis showed that self-stigma was an important factor negatively influencing adherence to treatment and significantly contributing to voluntary discontinuation of drugs. The level of self-stigma did not differ between diagnostic categories. Patients suffering from schizophrenia had the lowest adherence to treatment.

Conclusion: The study showed a significant correlation between self-stigma and adherence to treatment. High levels of self-stigma are associated with discontinuation of medications without a psychiatrist's recommendation. This connection was present in all diagnostic groups.

Keywords: self-stigma, adherence, discontinuation of drugs, mental disorders

Self-stigma and quality of life in patients with depressive disorder: a cross-sectional study

This article was published in the following Dove Press journal:
Neuropsychiatric Disease and Treatment
19 October 2016
[Number of times this article has been viewed](#)

Michaela Holubova^{1,2}
Jan Prasko¹
Marie Ociskova¹
Marketa Marackova¹
Ales Grambal¹
Milos Slepecky³

¹Department of Psychiatry, Faculty of Medicine and Dentistry, Palacky University Olomouc, University Hospital Olomouc, Olomouc,

²Department of Psychiatry, Hospital Liberec, Liberec, Czech Republic;

³Department of Psychology Sciences, Faculty of Social Science and Health Care, Constantine the Philosopher University, Nitra, Slovak Republic

Background: Self-stigma is a maladaptive psychosocial phenomenon that can affect many areas of patients' lives and have a negative impact on their quality of life (QoL). This study explored the association between self-stigma, QoL, demographic data, and the severity of symptoms in patients with depressive disorder.

Patients and methods: Patients who met the International Classification of Diseases, 10th revision, research criteria for depressive disorder were enrolled in this cross-sectional study. All outpatients completed the following measurements: the Quality of Life Enjoyment and Satisfaction Questionnaire, the Internalized Stigma of Mental Illness Scale, demographic questionnaire, and the objective and subjective Clinical Global Impression-Severity scales that measure the severity of disorder. A total of 81 depressive disorder patients (with persistent affective disorder – dysthymia, major depressive disorder, or recurrent depressive disorder) and 43 healthy controls participated in this study.

Results: Compared with the healthy control group, a lower QoL was observed in patients with depressive disorder. The level of self-stigma correlated positively with total symptom severity score and negatively with QoL. Multiple regression analysis revealed that the overall rating of objective symptom severity and score of self-stigma were significantly associated with QoL.


Conclusion: This study suggests a lower QoL in patients with depressive disorder in comparison with healthy controls and a negative impact of self-stigma level on QoL in patients suffering from depressive disorders.

Keywords: self-stigma, quality of life, depressive disorder

indicatori di self-stigma:

- si perde l'identità personale, sostituita da un senso del sé definito interamente dalla malattia
- tutte le attività della vita ruotano intorno all'essere paziente psichiatrico
- si sviluppa l'alienazione dagli altri
- si perdono i precedenti ruoli sociali
- si sviluppa un'idea di sé svalutata e socialmente indesiderata

Figure 2 A hierarchy of disclosure strategies



QUALITÀ DI VITA

«la percezione che la persona ha della sua posizione nella vita nel contesto dei sistemi culturale e valoriale in cui vive e in relazione ai suoi obiettivi, alle sue aspettative, ai suoi standard e alle sue preoccupazioni. Si tratta di un concetto ampio influenzato in modo complesso dalla salute fisica, dallo stato psicologico, dal livello di indipendenza, dalle relazioni interpersonali, dalle credenze personali e dalle loro interazioni con le caratteristiche salienti dell'ambiente»

(OMS, 1994)

QUALITÀ DI VITA CONNESSA ALLO STATO DI SALUTE

area dei sintomi fisici legati alla malattia e/o ai trattamenti: dolore, dispnea, perdita dei capelli




area della funzionalità fisica: capacità di svolgere le normali attività quotidiane come cura di sé, attività lavorativa e tempo libero

area dello stato emotivo: ansia, depressione, irritabilità

area delle relazioni sociali: interazioni con i familiari e con gli amici

... ma anche aspetti spirituali ed esistenziali, funzionalità sessuale, immagine corporea, soddisfazione per le cure ricevute e per la relazione medico-paziente...

CANCRO DEL DISTRETTO TESTA-COLLO



CANCRO DEL DISTRETTO TESTA-COLLO

Danker et al. (2010) – Social withdrawal after laryngectomy

- 40% dei 218 pz. con LT evita le conversazioni sociali
- 70% evita di partecipare ad attività sociali
- frequente la percezione di public stigma a causa del cambiamento della voce e della presenza di una tracheostomia

CANCRO DEL DISTRETTO TESTA-COLLO

PSYCHOSOCIAL ADJUSTMENT IN HEAD AND NECK CANCER: THE IMPACT OF DISFIGUREMENT, GENDER AND SOCIAL SUPPORT

Mark R. Katz, MD, FRCPC,¹ Jonathan C. Irish, MD, MSc, FRCSC,²
Gerald M. Devins, PhD, C. Psych,¹ Gary M. Rodin, MD, FRCPC,¹
Patrick J. Gullane, MB, FRCSC, FACS²

CANCRO DEL DISTRETTO TESTA-COLLO

1. danno estetico:

- importanza per l'immagine di sé
- importanza per le relazioni interpersonali
- importanza per la comunicazione
- altamente visibile
- come misurarla?

2. genere: le donne incontrano maggiori difficoltà nell'adattamento alle sequele del cancro?

3. supporto sociale: la percezione del supporto sociale pare correlata ad una migliore qualità di vita

CANCRO DEL DISTRETTO TESTA-COLLO

risultati:

1. livelli maggiori di depressione nelle donne e nei soggetti con più importante danno estetico
2. livelli maggiori di benessere psicologico in coloro che esperiscono maggiore supporto sociale
3. il supporto sociale modera gli effetti del danno estetico sul benessere psicologico nella donna, ma non nell'uomo
4. livelli più bassi di «life happiness» nelle donne e nei soggetti con più severo danno estetico

CANCRO DEL DISTRETTO TESTA-COLLO

conclusioni:

le donne sembrano più vulnerabili all'impatto del cancro del distretto testa-collo rispetto agli uomini e il loro adattamento dipende maggiormente dall'ambiente sociale

CANCRO DEL DISTRETTO TESTA-COLLO

THE IMPACT OF HEAD AND NECK CANCER AND FACIAL DISFIGUREMENT ON THE QUALITY OF LIFE OF PATIENTS AND THEIR PARTNERS

Lindsey E. Vickery, DClin Psychol,¹ Gary Latchford, PhD,² Jenny Hewison, PhD,²
Maggie Bellew, PhD,¹ Tricia Feber, Dip Nursing³

la variabile «danno estetico» ha una ridotta
ripercussione sulla qualità della vita

possibili interpretazioni:

1. il danno estetico è il prezzo da pagare per la sopravvivenza
2. danno estetico non così devastante
3. campione ridotto (chi non partecipa ha maggiori difficoltà di adattamento?)

CANCRO DEL DISTRETTO TESTA-COLLO

Acta Otolaryngol 2004; 124: 509–514

Taylor & Francis
healthsciences

Elderly Patients with Head and Neck Cancer: Physical, Social and Psychological Aspects after 1 Year

WYNIA DERKS¹, ROB DE LEEUW², JACQUES WINNUBST² and GERRIT JAN HORDIJK¹

From the ¹Department of Otorhinolaryngology and ²Research Group Psychology of Health and Illness, University Medical Center Utrecht, Utrecht, The Netherlands

Derks W, De Leeuw R, Winnubst J, Hordijk GJ. Elderly patients with head and neck cancer: physical, social and psychological aspects after 1 year. *Acta Otolaryngol* 2004 124: 509–514.

Objective—Elderly patients with head and neck cancer often receive non-standard treatment for reasons other than comorbidity. In this prospective study we investigated how elderly patients fare 1 year after treatment in comparison with their younger counterparts.

Material and Methods—Seventy patients aged 45–60 years and 51 patients aged ≥ 70 years with cancer of the oral cavity, pharynx (stage II–IV) or larynx (stage III–IV) participated in the study before treatment and 1 year later. Each patient was interviewed and given a questionnaire concerning physical functioning, social contacts, depressive symptoms, satisfaction with treatment and future expectations.

Results—At 12 months, patients in both age groups reported significantly more depressive symptoms and less social support than before treatment, and their Karnofsky Performance Score was lower. However, there were no differences between elderly and younger patients. Approximately 90% of those in both age groups said that they would choose the same treatment again, and there was no age difference regarding the impact of treatment or expectations for the future.

Conclusion—This study shows that the impact of treatment on quality of life did not differ between elderly and younger patients with head and neck cancer. Therefore, standard treatment should be considered in elderly patients if no severe contraindications exist. *Key words:* Depression, elderly, head and neck neoplasm, quality of life, social support therapy.

CANCRO DEL DISTRETTO TESTA-COLLO

QUALITY OF LIFE IN ELDERLY PATIENTS WITH HEAD AND NECK CANCER ONE YEAR AFTER DIAGNOSIS

Wynia Derks, MD,¹ Rob J. de Leeuw, PhD,² Gerrit Jan Hordijk, MD, PhD,¹
Jacques A. Winnubst, PhD²

¹ Department of Otorhinolaryngology, University Medical Center Utrecht, Department of Otorhinolaryngology, HP: G05.129, P.O. Box 85500, 3508 GA Utrecht, The Netherlands. E-mail: W.Derks@azu.nl

² Research Group Medical and Health Psychology, University Medical Center Utrecht, The Netherlands


Accepted 4 June 2004

Published online 30 September 2004 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/hed.20113

confermano i dati precedenti

CANCRO DEL DISTRETTO TESTA-COLLO

Acta Otolaryngol 2004 124: 231–236



REVIEW ARTICLE

Quality of Life Following Neck Dissections

SIMON N. ROGERS¹, ALFIO FERLITO², PHILLIP K. PELLITTERI³, ASHOK R. SHAHA⁴ and ALESSANDRA RINALDO²

From the ¹Department of Oral and Maxillofacial Surgery, University Hospital Aintree and Liverpool University Dental Hospital, Liverpool, UK, the Departments of Otolaryngology–Head and Neck Surgery, ²University of Udine, Udine, Italy and ³Geisinger Medical Center, Danville, Philadelphia, USA and the ⁴Head and Neck Service, Memorial Sloan–Kettering Cancer Center, New York, New York, USA

1. maggiore estensione della chirurgia del collo →
maggiore gravità della patologia della spalla
2. risparmio del nervo accessorio spinale
3. dolore maggiore per svuotamenti più estesi

CANCRO DEL DISTRETTO TESTA-COLLO

SHOULDER AND NECK MORBIDITY IN QUALITY OF LIFE AFTER SURGERY FOR HEAD AND NECK CANCER

C. P. van Wilgen, PT,^{1,2} P. U. Dijkstra, PT, MT, PhD,^{1,2} B. F. A. M. van der Laan, MD, PhD,³
J. Th. Plukker, MD, PhD,⁴ J. L. N. Roodenburg, DDS, PhD¹

¹ Department of Oral and Maxillofacial Surgery, University Hospital Groningen, PO Box 30.001, 9700 RB Groningen, The Netherlands. E-mail: c.p.van.wilgen@rev.azg.nl

² Department of Rehabilitation, University Hospital Groningen, Groningen, The Netherlands

³ Department of Otorhinolaryngology, Head & Neck Surgery, University Hospital Groningen, Groningen, The Netherlands

⁴ Department of Surgical Oncology, University Hospital Groningen, Groningen, The Netherlands

Accepted 27 February 2004

Published online 23 June 2004 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/hed.20052

riduzioni nell'abduzione e dolore localizzato alla spalla sono significativamente correlati all'esito in quattro domini:

1. funzionamento fisico
2. limitazioni dovute a problemi fisici
3. dolore
4. generale percezione della salute

CANCRO DEL DISTRETTO TESTA-COLLO

RADIATION-INDUCED XEROSTOMIA IN PATIENTS WITH HEAD AND NECK CANCER: PATHOGENESIS, IMPACT ON QUALITY OF LIFE, AND MANAGEMENT

Mark S. Chambers, DMD, MS,¹ Adam S. Garden, MD,² Merrill S. Kies, MD,³ Jack W. Martin, DDS, MS¹

¹ Department of Head and Neck Surgery, The University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Boulevard, Houston, Texas 77030-4009. E-mail: mchamber@mdanderson.org

² Department of Radiation Oncology, The University of Texas M. D. Anderson Cancer Center, Houston, Texas

³ Department of Head and Neck Medical Oncology, The University of Texas M. D. Anderson Cancer Center, Houston, Texas

Support Care Cancer (2008) 16:171–179

DOI 10.1007/s00520-007-0300-5

ORIGINAL ARTICLE

The influence of xerostomia after radiotherapy on quality of life

Results of a questionnaire in head and neck cancer

Piet Dirix · Sandra Nuyts · Vincent Vander Poorten ·
Pierre Delaere · Walter Van den Bogaert

la xerostomia è associata a:

1. dolore alla bocca
2. incremento delle carie dentali e delle infezioni orali
3. difficoltà nel parlare e nel deglutire
4. riduzione dell'alimentazione e perdita di peso

CANCRO DEL DISTRETTO TESTA-COLLO

QUALITY OF LIFE AS PREDICTOR OF WEIGHT LOSS IN PATIENTS WITH HEAD AND NECK CANCER

Karin M. Petruson, MD, Ewa M. Silander, RD, Eva B. Hammerlid, MD, PhD

From the Department of Otorhinolaryngology Head and Neck Surgery, Sahlgrenska University Hospital, S-413 45 Göteborg, Sweden. E-mail: karin.petruson@orlss.gu.se

Accepted 4 November 2004

Published online 14 February 2005 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/hed.20172

ORIGINAL ARTICLE

MALNUTRITION AND QUALITY OF LIFE IN PATIENTS TREATED FOR ORAL OR OROPHARYNGEAL CANCER

Harriët Jager-Wittenaar, RD,¹ Pieter U. Dijkstra, PT, MT, PhD,^{1,2,3} Arjan Vissink, DDS, MD, PhD,¹ Bernard F. A. M. van der Laan, MD, PhD,⁴ Rob P. van Oort, DDS, PhD,¹ Jan L. N. Roodenburg, DDS, MD, PhD¹

¹ Department of Oral and Maxillofacial Surgery, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands. E-mail: h.jager@onco.umcg.nl

² Department of Rehabilitation, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

³ Graduate School for Health Research, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

⁴ Department of Otorhinolaryngology-Head and Neck Surgery, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

Accepted 31 March 2010

Published online 7 September 2010 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/hed.21473

- perdita di peso (incidenza 35-50%) → incrementa morbilità e mortalità e riduce la tolleranza alla radioterapia e alla chemioterapia
- una perdita di peso che supera il 10% del peso in 6 mesi → malnutrizione e peggiore QOL anche a 3 anni

CANCRO DEL DISTRETTO TESTA-COLLO

QUALITY OF LIFE AND ILLNESS PERCEPTIONS IN PATIENTS WITH RECENTLY DIAGNOSED HEAD AND NECK CANCER

Margreet Scharloo, PhD,¹ Robert J. Baatenburg de Jong, MD, PhD,² Ton P. M. Langeveld, MD, PhD,² Els van Velzen-Verkaik, RN,² Margreet M. Doorn-op den Akker, MSW,² Adrian A. Kaptein, PhD¹

¹ Psychology Unit, Leiden University Medical Center, PO Box 9555, 2300 RB Leiden, The Netherlands.
E-mail: m.scharloo@lumc.nl

² Department of Otolaryngology and Head and Neck Surgery, Leiden University Medical Center, Leiden, The Netherlands

Apparently, it is not so much the disease itself but the way the patient perceives and responds to the disease that determines an individual's QOL.^{6,9}

Eliciting the patient's views of his or her illness offers the potential for a better understanding of (mal)adaptive response to illness and treatment and has implications for medical practice.

conclusioni:

pazienti con percezioni di malattia più negative prima del trattamento (aumentata attenzione ai sintomi, aumentata credenza rispetto alla possibilità di una recidiva, maggiore tendenza ad autocolpevolizzarsi e reazioni emozionali più forti alla malattia) hanno una più ridotta qualità di vita

CANCRO DEL DISTRETTO TESTA-COLLO

Patients' views on the utility of quality of life questionnaires in head and neck cancer: a randomised trial

Mehanna, H.M.,*[†] & Morton, R.P.[‡]

**Department of Head and Neck Surgery, University Hospitals Coventry and Warwickshire, Coventry, [†]Heart of England Foundation Trust, Birmingham, UK, and [‡]Department of Otorhinolaryngology, Head and Neck Surgery, Auckland City Hospital, Auckland, New Zealand*

Accepted 1 May 2006
Clin. Otolaryngol. 2006, 31, 310-316

Objectives: (i) To evaluate head and neck cancer patient perspectives regarding the usefulness of quality of life questionnaires in communicating their health problems to clinicians and (ii) to identify the quality of life questionnaire that head and neck cancer patients find most useful.

Design: Randomised questionnaire study. Patients completed all four validated head and neck cancer quality of life questionnaires – European Organisation for Research and Treatment of Cancer (EORTC), Functional Assessment of Cancer Therapy Scale (FACT) HN35, Washington quality of life questionnaire, Auckland quality of life questionnaire. Order of questionnaire presentation was randomised to counterbalance order effects.

Setting: Tertiary referral head and neck cancer centre.
Participants: Eighty patients diagnosed and treated for head and neck cancer. Exclusion criteria: blindness, learning difficulties or inability to understand or read English.

Main outcome measures: Patient ratings of perceived usefulness and preferences of studied questionnaires.

Results: Patients reported high relevance to their problems and high ease of understanding of all questionnaires, with FACT scoring highest (79% and 89%, respectively); 58% of participants (67% respondents) would like to complete a questionnaire in clinic, as it would help them describe their health problems to their doctors; 28% of participants did not. Almost half preferred a particular quality of life questionnaire, FACT being most preferred. Length of questionnaire did not affect reported usefulness, but most would prefer a short questionnaire (<20 items).

Conclusions: Patients report that head and neck cancer quality of life questionnaires effectively describe their health concerns. Most are in favour of completing quality of life questionnaires in clinic, as an aid for describing health problems to clinicians. There appears to be a difference between clinicians and patients regarding the perceived usefulness of quality of life questionnaires in the clinic setting, which needs to be highlighted to clinicians.

CANCRO DEL DISTRETTO TESTA-COLLO

Otolaryngology–Head and Neck Surgery (2007) 136, 405-410

ORIGINAL RESEARCH

Quality-of-life impact of participation in a head and neck cancer support group

Kalpesh T. Vakharia, MD, M. Jafer Ali, MD, and Steven J. Wang, MD,
San Francisco, CA



OBJECTIVE: To assess if participation by patients in a head and neck cancer support group improves perceived quality of life (QOL).

STUDY DESIGN AND SETTING: Subjects for this study included 47 patients at a tertiary Veterans Affairs Medical Center who were previously treated for head and neck cancer. This was a quasi-experimental, post-test study comparing the QOL of 24 patients who participated in a head and neck cancer support group with 23 patients who did not participate. The validated University of Michigan Head and Neck Quality of Life (HNQOL) instrument was used to evaluate head and neck cancer-related QOL.

RESULTS: Patients who participated in the head and neck cancer support group exhibited significantly better scores in the domains of eating, emotion, and pain as well as in the global bother and response to treatment questions of the HNQOL instrument compared with those patients who did not participate. Additional subgroup analysis comparing age, type of treatment, and length of time since cancer diagnosis suggests that these variables were less important predictors of QOL than was support group participation.

CONCLUSIONS: Our findings suggest that patient participation in a head and neck cancer support group is associated with improved QOL.

CEREBROLESIONE



CEREBROLESIONE

DISABILITY AND REHABILITATION, 2002; VOL. 24, NO. 5, 259–265



Predictors of quality of life following stroke

A. E. MACKENZIE* and A. M. CHANG

Department of Nursing, Floor 7 CCV Chung Chi, The Chinese University of Hong Kong,
Shatin, NT, Hong Kong

- durata del ricovero
- ictus precedenti
- livello di funzionamento
- supporto sociale

CEREBROLESIONE

*Acta Neurol Scand 2003; 107: 324-329
Printed in UK. All rights reserved*

Copyright © Blackwell Munksgaard 2003

ACTA NEUROLOGICA
SCANDINAVICA
ISSN 0901-6114

Quality of life in stroke patients

Jaracz K, Kozubski W. Quality of life in stroke patients.
Acta Neurol Scand 2003; 107: 324-329. © Blackwell Munksgaard 2003.

Objectives – To describe global and domain-specific quality of life (QOL) after stroke and to identify the factors that are important for post-stroke QOL. *Material and methods* – A hospital-based sample of 72 stroke patients was followed up for 6 months after stroke onset. QOL was assessed using the Polish version of the Quality of Life Index. Regression analysis was performed to identify the variables that best predicted QOL. *Results* – The overall QOL of stroke patients was relatively good, although worse than that of subjects in a comparison group. The highest QOL was found in the ‘Family’ domain, and the lowest in the ‘Health and functioning’ domain. Emotional support, depression and functional disability were three separate variables explaining 38% of the variance in QOL. *Conclusions* – Strengthening of family support, treatment of depression and reduction of physical dependence may be the decisive factors in improving post-stroke QOL.

K. Jaracz¹, W. Kozubski²

Departments of ¹Nursing and ²Neurology, University of Medical Sciences, Poznań, Poland

Key words: depression; quality of life; social support; stroke

Krzyszyna Jaracz, Department of Nursing, University of Medical Sciences, Dabrowskiego 79, Poznań, Poland

Tel: +01033 61 847 74 91 ex. 189

Fax: +01033 61 847 74 90

e-mail: jkrzys@poczta.onet.pl

Accepted for publication August 2, 2002

CEREBROLESIONE



Disponibile en ligne sur

 www.sciencedirect.com

Elsevier Masson France

 www.em-consulte.com/en



Annals of Physical and Rehabilitation Medicine 54 (2011) 376–390

Original article / Article original

Assessment of quality of life in stroke patients with hemiplegia

Évaluation de la qualité de la vie après un accident vasculaire cérébral

K. Laurent^{a,b}, M.-P. De Sèze^{a,b}, C. Delleci^{a,b}, M. Koleck^c, P. Dehail^{a,b},
 J.-M. Orgogozo^d, J.-M. Mazaux^{a,b,*}

Abstract

Stroke is a major public health issue. Even though most hemiplegic stroke patients may obtain a good functional outcome, many remain dissatisfied with their lives. Indeed, quality of life and subjective well-being should be taken into account in any assessment of stroke survival. *Objective.* – To assess long-term quality of life in stroke patients (compared with healthy controls) and the corresponding determinants and predictive factors.


Method. – The patient population consisted of 80 of the 217 first-stroke survivors treated between January and June 2005 in the Clinical Neurosciences Department at Bordeaux University Hospital. After a mean follow-up period of 2 years, 24 patients were interviewed in their homes and data from the 56 others were obtained in a telephone interview. Demographic information, clinical status on admission and functional status (as assessed by Barthel Index) and depression (on the ADRS) at the time of the study visit were recorded. Quality of life was assessed by using the Sickness Impact Profile (SIP-65) and Brånholm and Fugl-Meyer's Satisfaction with Life Scale (LiSat 11). The patients' data were compared with those from 149 healthy controls.

Results. – Life satisfaction and quality of life were significantly impaired in stroke patients, compared with controls. All life domains were impaired. The worst scores were observed for independence and health-related items in the LiSat 11 and the physical and communication items in the SIP-65. Quality of life was strongly correlated with functional independence, the persistence of hemiplegia and depressive mood, which is in agreement with literature findings. Neither gender nor the initial Rankin score had a significant impact on these parameters.

Discussion–Conclusion. – Quality of life at 2 years is significantly impaired in stroke survivors and seems more difficult to predict than functional independence. However, in addition to these objective results, our interviews suggest that receiving adequate social support might be as important to patients as recovering independence.

CEREBROLESIONE

Brain Injury, 20 August 2005; 19(9): 657–665



ORIGINAL PAPER

Self-concept can be defined as a collection of representations reflecting a person's beliefs about his/her own functioning in various life dimensions, such as physical self-concept and social self-concept

Self-concept and quality of life following acquired brain injury: A pilot investigation

C. D. VICKERY¹, S. T. GONTKOVSKY¹, & J. S. CAROSELLI²

¹*Methodist Rehabilitation Center, Jackson, MS, USA* and ²*The Institute for Rehabilitation and Research, Houston, TX, USA*

(Received 7 January 2004; accepted 12 July 2004)

Abstract

Objective: To investigate through pilot exploration the relationships between depression, self-concept and perceived quality of life (QoL) in post-acute patients with acquired brain injury (ABI).

Methods: Nineteen patients with ABI were administered the Beck Depression Inventory-II and the Quality of Life Inventory, along with the Tennessee Self-Concept Scale-2 and the Head Injury Semantic Differential Scale, measures of self-concept. The relationships between these measures were explored using correlational analyses.

Results: Ratings of self-concept were correlated with perceived QoL, suggesting that poorer view of self was associated with lower subjective QoL. Additionally, depressive symptoms were associated with lower QoL ratings, consistent with previous research.

Conclusions: These results suggest that intra-personal variables, such as self-concept and depression, impact the perceived QoL of the ABI survivor. Future research exploring the mediating effects of these variables on QoL may clarify this relationship and may aid in developing more effective interventions for these individuals.

CEREBROLESIONE

Brain Injury, June 2005; 19(6): 403–415



ORIGINAL ARTICLE

Female body image following acquired brain injury

H. F. R. HOWES, S. EDWARDS, & D. BENTON

University of Wales, Swansea, Wales, UK

(Received 5 February 2004; accepted 8 November 2004)

Abstract

Primary objective: To investigate body image concerns and psycho-emotional health in females with acquired brain injury (ABI).

Research design: Between subjects design.

Methods and procedures: Thirteen females with ABI and thirteen matched controls were assessed on a number of variables relating to body image, self-esteem and affect.

Main outcomes and results: Indicated that women with ABI had a different profile of body image dissatisfaction than controls and that in some areas it was significantly worse. The women with ABI had significantly lower self-esteem and higher levels of depression than controls and this related to body image.

Conclusion: Significant changes in body image and self-concept are potentially damaging to quality of life post-injury and suggest a role for targeted therapeutic work with women with ABI.

STRABISMO

scopo principale del trattamento → allineare gli assi visivi in modo da raggiungere un'unica visione binoculare con una buona fusione delle immagini

altri vantaggi:

- miglioramento delle alterazioni posturali del capo
- ampliamento e centralizzazione del campo visivo
- miglioramento della motilità oculare
- miglioramento dello sviluppo psicomotorio
- recupero di un «normale» aspetto fisico

STRABISMO

problematiche psicosociali ed emotive tipiche:

- ridotta autostima
- pregiudizio sociale
- bullismo
- incremento dell'ansia sociale
- relazioni interpersonali fragili
- problematiche nel trovare un lavoro

STRABISMO

Quality of life of patients with strabismus

Qualidade de vida em pacientes estrábicos

GERALDO DE BARROS RIBEIRO¹, ANA GABRIELA ZUM BACH¹, CAMILA MAIA FARIA¹, SUZE ANASTÁSIA¹, HENDERSON CELESTINO DE ALMEIDA¹

ABSTRACT

Purpose: To evaluate the impact of strabismus on quality of life.

Methods: This cross-sectional study included 101 individuals aged 7-67 years with strabismus. They were interviewed and made to answer a questionnaire with 20 questions intended to assess the individual's interaction with their social and working environment, sensory perception, and limitations in their daily lives. There were five types of possible responses for each question: never, rarely or very few times, sometimes, frequently, and always. The questionnaire was based on the AS-20 and contained 11 questions assessing psychosocial aspects and 9 questions assessing functional aspects. Among those who were interviewed, 24.8% had undergone surgical correction of strabismus.

Results: We observed a significant impact of strabismus on the quality of life of the interviewed individuals. Feelings of sadness and inferiority because of strabismus were reported by 74.2% and 58.4% respondents, respectively. In terms of functionality, 12.1% reported difficulty in reading, 14% said they had difficulty in depth perception (stereopsis), and 17.8% frequently or always associated pain or burning sensation in the eyes to strabismus. A significant difference was detected in the quality of life scores for the psychosocial aspect among patients who had and had not undergone surgery (Wilcoxon test, 158; $p < 0.001$). Individuals who had undergone surgery had a better quality of life from the psychosocial perspective.

Conclusions: In this evaluation, we found a significant negative interference of strabismus on quality of life from both the functional and psychosocial perspectives. This demonstrated the importance of treatment for strabismus, regardless of age, because it can interfere with the functional well-being of the individual.

RESUMO

Objetivo: Avaliar a interferência do estrabismo na qualidade de vida em indivíduos estrábicos.

Métodos: Estudo transversal envolvendo 101 indivíduos estrábicos entre 7 e 67 anos. Eles foram entrevistados e responderam um questionário com 20 perguntas aplicadas pelos pesquisadores. As perguntas tinham como objetivo avaliar a interação do indivíduo com o seu meio social, de trabalho, percepção sensorial e limitações na sua vida diária, com cinco tipos de respostas para cada pergunta: nunca, raramente ou muito pouco, algumas vezes, frequentemente e sempre. O questionário foi baseado no AS-20 e contendo 11 questões avaliando aspectos psicossociais e 9 questões avaliando aspectos funcionais. Entre os indivíduos entrevistados, 24,8% haviam sido submetidos a correção cirúrgica do estrabismo.

Resultados: Observou-se impacto significativo do estrabismo sobre a qualidade de vida dos indivíduos avaliados. Sentir-se incomodado ou inferiorizado em decorrência do estrabismo foram relatados por 74,2% e 58,4% dos entrevistados respectivamente. No quesito de funcionalidade, 12,1% relataram dificuldade para ler, 14% disseram ter dificuldade na percepção de profundidade (estereopsia) e 17,8% associaram dor ou ardor nos olhos ao estrabismo frequentemente ou sempre. Diferença significativa foi detectada com respeito ao escore de qualidade de vida no aspecto psicossocial entre indivíduos que realizaram e não realizaram cirurgia (teste de Wilcoxon = 158, p -valor $< 0,001$). Indivíduos que realizaram cirurgia possuíam melhor qualidade de vida no aspecto psicossocial.

Conclusões: Nessa avaliação, percebe-se uma interferência negativa muito significativa do estrabismo na qualidade de vida, tanto funcional como psicossocial nos indivíduos pesquisados. Isso demonstra a importância do tratamento do estrabismo, independente da idade, pois ele pode interferir muito na qualidade de vida e bem estar psicossocial e funcional do indivíduo.

STRABISMO

Annex. Questionnaire on the impact of strabismus on patient quality of life

Questions	Never	Rarely or very few times	Sometimes	Frequently	Always
1- Do you notice you have strabismus (crossed eyes or lazy eyes)?					
2- Does strabismus bother you?					
3- When you interact with other people, do you feel they notice your strabismus?					
4- Do you feel different (inferior) because of strabismus?					
5- Do you feel that strabismus hinders your performance (at school or at work)?					
6- Do you feel that strabismus affects your relations with other people?					
7- Do you feel that your opportunities (social interaction or at work) are fewer because of strabismus?					
8- Do you imagine what other people are thinking about your eyes?					
9- Do you have more difficulty in making friends because of strabismus?					
10- Do you feel uncomfortable if someone asks something about your eyes?					
11- Do you feel that people avoid looking at you because of strabismus?					
12- Does strabismus hinder your vision?					
13- Do you have difficulty in reading because of strabismus?					
14- Do you feel more stressed out (anxious) because of strabismus?					
15- Do you have difficulty in depth perception (assess the distance of objects)?					
16- Do you feel any discomfort such as pain or burning sensation in the eyes because of strabismus?					
17- Do you have to close one of your eyes to read?					
18- Do you have diplopia (double vision)?					
19- Do you feel eye strain because of strabismus?					
20- Do you think about your strabismus so frequently that it hinders your concentration?					

STRABISMO

Lo strabismo rappresenta un problema?

49.5% sempre

Ti senti inferiore a causa dello strabismo?

41.5% sempre

Lo strabismo ti crea dei problemi sul lavoro o a scuola?

22.7% sì

Lo strabismo ti ha mai creato problemi nei rapporti interpersonali?

53% sì

Le persone evitano di guardarti a causa dello strabismo?

16.8% spesso + sempre

STRABISMO

Changes in Health-Related Quality of Life 1 Year Following Strabismus Surgery (Am J Ophthalmol 2012;153: 614–619)

SARAH R. HATT, DAVID A. LESKE, LAURA LIEBERMANN, AND JONATHAN M. HOLMES

- **PURPOSE:** To report changes in health-related quality of life (HRQOL) 1 year following strabismus surgery.
- **CONCLUSIONS:** Adult strabismus patients who remain successfully aligned show continued improvement in both psychosocial and function scores from 6 weeks to 1 year postoperatively, indicating that improvement in HRQOL is long-lasting. Evaluation of HRQOL should be considered when reporting strabismus surgery outcomes in clinical trials, and may prove helpful in assessing outcomes in clinical practice.

PSICOPATOLOGIA

Acta Psychiatr Scand 2006; 113 (Suppl. 429): 46-50
 All rights reserved
 DOI: 10.1111/j.1600-0447.2005.00717.x

Copyright © 2006 Blackwell Munksgaard
 ACTA PSYCHIATRICA
 SCANDINAVICA

Determinants of quality of life in people with severe mental illness

Hansson L. Determinants of quality of life in people with severe mental illness.

Acta Psychiatr Scand 2006; 113 (Suppl. 429): 46-50. © 2006 Blackwell Munksgaard.

Objective: The objective of this article was to review some methodological issues in this field and give an overview of empirical research findings with a special focus on factors associated with or affecting subjective quality of life in people with a severe mental illness.

Method: A selective review of relevant scientific literature on quality of life in severe mental illness was conducted.

Results: Subjective quality of life in people with a severe mental illness is only to a lesser extent related to external life conditions. Major determinants are psychopathology, especially symptoms of depression and anxiety, and aspects of the social network. Personality related factors such as self-esteem are also influential. Comparative studies have further shown that patients in community care settings have a better subjective quality of life than patients in hospital settings.

Conclusion: Efforts to improve subjective quality of life in people with severe mental illness should include a careful monitoring of depressive and anxiety symptoms, and pay particular attention to assessment of and interventions against unmet needs. Further, such interventions should stress a strengthening of the social support of the clients. It is also important to pay attention to mediators of changes in subjective quality of life such as self-esteem, mastery, autonomy, and self-efficacy.

L. Hansson

Department of Health Sciences, Division of Nursing
 Lund University, Lund, Sweden

Key words: subjective quality of life; psychopathology; social support; personality; treatment outcome

Lars Hansson, Department of Health Sciences, Division of Nursing, Lund University, PO Box 157, SE-221 00 Lund, Sweden.

E-mail: lars.hansson@omv.luse

This paper was read in a preliminary version at the 3rd International Zurich Conference on Clinical and Social Psychiatry, Zurich, 25-27 September, 2003. The conference and publication of this supplement were exclusively sponsored by Eli Lilly, Suisse.