



# Neanderthal language revisited: not only us

Dan Dediu<sup>1,2,4</sup> and Stephen C Levinson<sup>3,4</sup>

Here we re-evaluate our 2013 paper on the antiquity of language (Dediu and Levinson, 2013) in the light of a surge of new information on human evolution in the last half million years. Although new genetic data suggest the existence of some cognitive differences between Neanderthals and modern humans — fully expected after hundreds of thousands of years of partially separate evolution, overall our claims that Neanderthals were fully articulate beings and that language evolution was gradual are further substantiated by the wealth of new genetic, paleontological and archeological evidence briefly reviewed here.

## Addresses

<sup>1</sup> Language and Genetics Department, Max Planck Institute for Psycholinguistics, Wundtlaan 1, 6525 XD Nijmegen, The Netherlands

<sup>2</sup> Collegium de Lyon, Institut d'Études Avancées, 24 rue Baldassini, 69007 Lyon, France

<sup>3</sup> Language and Cognition Department, Max Planck Institute for Psycholinguistics, Wundtlaan 1, 6525 XD Nijmegen, The Netherlands

<sup>4</sup> Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, Netherlands

Corresponding authors: Dediu, Dan ([Dan.Dediu@mpi.nl](mailto:Dan.Dediu@mpi.nl)), Levinson, Stephen C ([Stephen.Levinson@mpi.nl](mailto:Stephen.Levinson@mpi.nl))

Current Opinion in Behavioral Sciences 2018, 21:49–55

This review comes from a themed issue on **The evolution of language**

Edited by **Christopher Petkov** and **William Marslen-Wilson**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 3rd February 2018

<https://doi.org/10.1016/j.cobeha.2018.01.001>

2352-1546/© 2018 Elsevier Ltd. All rights reserved.

## Introduction


In 2013 we published a paper [1] arguing that vocal language has had a long and gradual evolution in the 1.5 my since *Homo erectus*, and that Neanderthals were articulate humans, not very different from us. We marshaled paleontological, archeological and genetic evidence against the saltationist view that language evolved abruptly within the last 100 ky (e.g. [2]). Since then there has been a flurry of new evidence from paleontology, archeology and ancient DNA that, we argue here, largely supports our views of a much deeper antiquity and gradualism.

## The human tree

One striking finding is that many human lineages were coexisting just 300 kya (including *Homo naledi*, *Homo floresiensis*, Denisovans, Neanderthals and even anatomically modern humans [3]), another being the degree of reticulation in the human tree evidenced by genetics. Recent data suggests that at least 50 ky of interactions on a vast geographic scale from the Levant to Siberia and Western Europe gave plenty of scope for interbreeding; [Figure 1](#) shows some of the gene flow across the Neanderthals, Denisovans and ourselves.

The ancestors of Neanderthals and Denisovans left Africa c. 650 kya, and by 450 kya had diverged into these two lineages, and there are indications of early Neanderthal and Denisovan interbreeding [4], and of interbreeding between Denisovans and yet another archaic hominin. The Neanderthals then interbred with anatomically modern humans (AMH) on at least three occasions, likely when AMH first came into the Levant 100–120 kya [5,6], and also during AMH's move into glacial Europe (c. 50 kya [7]). For example, a Siberian Neanderthal has AMH genes introgressed c. 110 kya [7], and an early AMH with Neanderthal anatomical features c. 40 kya from Romania [8] had Neanderthal ancestry just 4–6 generations before [9]. Recent data suggests that while Neanderthals and Denisovans clearly form a clade, the mtDNA groups the former with AMH due to their original mtDNA being replaced by an AMH-like lineage probably through interbreeding before c. 270 kya [10]. It is now clear that such interbreeding did leave traces in contemporary modern humans outside Africa at the level of a few percent of the genome, with regional and inter-individual variation probably due to repeated interbreeding with Neanderthals and Denisovans [5,11,12]. Contact between human lineages is also revealed by other lines of evidence including the oral commensal microbe *Methanobrevibacter oralis* from a Spanish Neanderthal pointing to his ancestors having had contact with an African strain presumably from the AMH in the Levant c. 126 kya [13], and the sexually transmitted oncogenic *human papillomavirus 16* which shows strains having split c. 450 kya mirroring the divergence between Neanderthals and AMH, but later with the Neanderthal strain re-infecting AMH after c. 120 kya [14]. Leaving aside the functional relevance of these genomic introgressions for now, what they overwhelmingly highlight is the interwoven histories of the human lineages within and outside Africa during the last half million years.

Figure 1



Schematic representation of human evolution (based on [15]) focusing on the lineages leading to ourselves (blue), Neanderthals (green) and Denisovans (magenta) and highlighting probable interbreeding (numbered red arrows). 1: 2.5–5.8% Denisova genome from archaic hominin having diverged 0.9–1.4 mya [16]; 2: mtDNA introgressed c. 270 kya into a Neanderthal (Hohlenstein-Stadel, Germany) from an African lineage leading or related to AMH [10] (see also [4]); 3: at least 0.5% genome coming from a Neanderthal population closer related to the Altai Neanderthals [16]; 4: 1.0–7.1% gene flow from AMH into Altai Neanderthals [7] (see also [9]); 5,6,8,9: multiple introgressions from Neanderthals into various modern human populations outside Africa resulting in about 2% (regionally and inter-individually variable, slightly more in East Asia) Neanderthal DNA [5<sup>\*\*</sup>,11<sup>\*\*</sup>,12<sup>\*\*</sup>]; 7: Denisova introgression resulting in about 2–4% Denisovan DNA in Melanesia (less in e.g. South Asia [5<sup>\*\*</sup>,12<sup>\*\*</sup>]).

### Functional genetic differences between AMH and Neanderthals

With almost 20 partial or full Neanderthal genomes in hand, the differences between ‘us’ and ‘them’ are quantitatively very modest indeed (e.g. [16] page 48 state that

there are ‘96 fixed amino acid substitutions in a total of 87 proteins and in the order of three thousand fixed changes that potentially influence gene expression’), but they might potentially have high functional relevance and complex, often indirect and still poorly understood

phenotypic consequences. For example, some of these genes with fixed changes are expressed in the developing neocortex [16], other are involved in skin, hair and bone structure, while others affect the immune and reproductive systems [11,17<sup>••</sup>]. Nevertheless, it is now possible to explore the function and evolutionary fate of archaic introgressions in a ‘fossil-free’ manner by examining the introgressions in modern populations [17].

For understanding language evolution in particular, the genes involved in brain development and the vocal organs are of course especially interesting but, unfortunately, linking molecular genetics to language and speech is an extremely complex endeavor [18,19]. Thus, there has been only slow progress in understanding the functional implications of the non-coding differences between modern and archaic humans in the ‘star’ gene *FOXP2* (such as the *POU3F2* putative binding site in intron 8 [20]), but it seems likely that non-coding changes in this and other language-implicated genes such as *ROBO1*, *ROBO2*, and *CNTNAP2* rose to high frequency after the separation of AMH from archaic hominins [21]. However, these genes are expressed not only in the brain, increasing the difficulty of attributing these putative selective pressures to language and speech.

Another approach is to locate sequences of archaic DNA that have increased in frequency or, more often, that have been systematically purged from the modern genome, and try to understand their functional implications. These ‘genomic deserts’ from which archaic DNA seems to have been eliminated [5<sup>••</sup>,12<sup>••</sup>,22<sup>••</sup>] are (mostly non-coding) regions located preferentially near protein-coding genes [22<sup>••</sup>], evolutionarily conserved regions [12<sup>••</sup>,23<sup>•</sup>,24], genes involved in brain development [5<sup>••</sup>], including *FOXP2* [5<sup>••</sup>,12<sup>••</sup>], and genes expressed in the testes and in meiotic germ cells and potentially relevant for male hybrid sterility [11<sup>••</sup>,12<sup>••</sup>,25<sup>••</sup>]. The most plausible explanation for this loss of archaic DNA is negative selection [5<sup>••</sup>,12<sup>••</sup>], an inference reinforced by other lines of evidence including the tissue-specific (especially in the brain and testes) downregulation of introgressed archaic DNA [26<sup>••</sup>] and the association of some archaic sequences with various pathologies [27<sup>••</sup>] in existing humans, possibly resulting from a combination of genomic incompatibilities [5<sup>••</sup>,12<sup>••</sup>] and many weakly deleterious alleles becoming visible to purifying selection after introgression in the larger AMH population [23<sup>•</sup>,28<sup>•</sup>]. However, the pattern is not all about loss: there are regions of archaic DNA that have dramatically increased in frequency, especially near genes involved in immunity [5<sup>••</sup>,29–33], metabolism [5<sup>••</sup>,12<sup>••</sup>] and skin, hair and bone phenotypes [5<sup>••</sup>,11<sup>••</sup>,32,34], pointing to advantageous adaptations developed by the other hominins in their non-African environments [32,33,35<sup>••</sup>]. In understanding these patterns, it is important to remember that Neanderthals were a heavily inbred population (the Altai

individual’s parents were related as half-siblings) with very low effective population sizes (possibly by an order of magnitude lower than the early AMH’s), which may have lowered their fitness by as much as 40% [7,23<sup>•</sup>,28<sup>•</sup>].

One interesting development has been the exploration of the genes involved in the reduction of facial prognathism, dentition and jaw, and the correlated globularization of the brain, a process that extends over 2 my of human evolution and that has traditionally been viewed as related to changes in diet, cooking and technology. However, the last stages of the process, and the transition from Neanderthal to AMH skull-shape (more globular), might have been driven by language. Some authors argue that globularization may have originated in ‘self-domestication’ with associated neotenization, affording better neural connectivity, and they highlight wide networks of genes potentially involved in this process [36–39]. However, the recent reanalysis of 300 ky early AMH skulls from Morocco shows that these skulls do not have a fully globular shape despite their fully modern faces [3<sup>•</sup>,40<sup>•</sup>]. Other researchers argue that facial reduction may have been driven by speech, more precisely by the optimization of the 1:1 oral versus supralaryngeal vocal tracts proposed to be crucial for speech production [41,42]. Exciting recent comparisons of methylation patterns between modern and archaic humans ([43], D Gokhman *et al.*, <https://www.biorxiv.org/content/early/2017/10/03/106955>) found that modern human-specific changes preferentially affect the face, the vocal tract (especially the larynx) and the pelvic region, suggesting that there were further changes in our lineage, after the split from the Neanderthals, possibly affecting vocal tract anatomy and physiology. Nevertheless, even if these changes did affect (or even evolved in response to) speech production, they suggest that probably our lineage tweaked an already-existing speech production apparatus instead of suddenly evolving a new one from scratch.

## Neanderthal cognition as evidenced by archeology

Neanderthals once covered a vast and climatically and ecologically diverse range, from Siberia to Gibraltar, and contrary to earlier stereotypes, they did adapt their diet and technology to local conditions [13,44]. In northern latitudes, they were predominantly carnivorous, had advanced leather processing technology (using *lissoirs*, probably later borrowed by AMH [45]), with modeling studies suggesting advanced winter clothing [46] using stone and bone awls (such as found in the Grotte du Renne) as sewing aids [47], but in general their diet was apparently very flexible and adapted to the local environment, including both meat and vegetable foods [48] to varying proportions [13,49–51]. Bitter medicinal herbs were consumed [52], a Neanderthal with dental abscess appears to have treated himself with poplar bark (a source of salicylic acid) and possibly mold-produced antibiotics

[13], and there are indications of possible dental manipulations [53] — all suggesting medicinal expertise.

The purported differences in technology between AMH and Neanderthals now appear much smaller following the clear association of the *Chatelpéronian* at the Grotte du Renne with the Neanderthals [54<sup>\*</sup>], new analyses suggesting that there were no differences in raw material efficiency [55], that bone was used for tools [56], that hafting might go back to 240–270 kya [57] and that manganese dioxide was used to produce fire on demand 50 kya [58], and only after c. 40 kya (as the Neanderthals were disappearing) did AMH technology become clearly superior.

One critical question has been whether there are any traces of Neanderthal symbolic activity, with earlier potential examples often been dismissed as excavation or dating errors. However, the unequivocal recovery of Neanderthal DNA at Grotte du Renne [54<sup>\*</sup>] seems to connect them, the advanced *Chatelpéronian* technology, and the elaborate necklaces of animal teeth and shells found there (but see [59]). Recently, a cave in Spain, apparently used as a mortuary site involving fire and hunting trophies but not inhabited, has yielded Mousterian layers containing numerous small hearths with over 30 antlers and horn-cores associated with an infant burial (E Baquedano *et al.*, Session 7, ESHE 2016, Madrid). There are suggestions that Neanderthals at several sites may have used long bird feathers for personal adornment [60], but probably the most spectacular recent findings are the c. 40 kya decorated raven wing bone from Crimea [61] and the c. 170 kya circular constructions from broken stalagmites (collectively weighting more than two tones) more than 300 m deep in the Bruniquel cave [62], for which it is hard to imagine any reason other than ceremonial.

### Inferences concerning language and speech

In what follows we adopt the view that language involves a cognitive architecture that maps sounds (or gestures) into meaning through a series of combinatorial structures, while insisting that language is also a motoric and perceptual skill. Due to the fact that existing languages vary widely in their organization, it is not possible to be very precise about the general properties of those structures, except that they involve a series of transductions across levels of representation (e.g. phonetics, phonology, morphology, syntax, semantics, pragmatics) without 1:1 mappings. Not all languages, for example, exhibit phrase structure at a surface level [63], but developed languages probably all involve hierarchical and recursive structure if not at a syntactic level, then at a semantic and pragmatic one. The critical function is to render propositional content and associated illocutionary force into a perceptual medium, with content recoverable with the aid of contextual inference. Although the inference of prehistoric linguistic structure will always be indirect, the fact that advanced technology itself involves hierarchical and

recursive structure (as in the manufacture of a composite tool) provides circumstantial evidence for the existence of the planning, memory and motor skills also involved in language [64]. Language affords culture-carrying capacity (e.g. there are no advanced technologies without language), and this linkage allows reasonable inferences from the archeological record [65]. Therefore, we think it is overwhelmingly likely that Neanderthals were as much articulate beings as we ourselves are, that is, with large vocabularies and combinatorial structures that allowed propositional content and illocutionary force to be conveyed. Only such an advanced communication system could have carried the advanced cultural adaptations that Neanderthals exhibited. However, some of the new findings, especially those coming from the genetics of brain development and neurocognitive disorders, seem to strongly suggest that, despite a profound shared humanity, archaic and modern humans were not identical [66].

The arguments for speech given in our 2013 paper still hold: Neanderthals had the modern (or near-modern) vocal apparatus ([43], D Gokhman *et al.*, <https://www.biorxiv.org/content/early/2017/10/03/106955>), breathing control, and acoustic sensitivity (audiograms) involved in modern speech. Since then, there is better evidence that the Neanderthal cochlear volume and audition were similar to modern humans and different from the chimpanzees [67], that despite differences in the morphology of the ear ossicles (probably due to differences in cranium) they were functionally very similar probably due to selective pressures for the maintenance of the same auditory profile [68<sup>\*</sup>], while the biomechanical modeling of the Kebara 2 Neanderthal hyoid showed that it is extremely similar to the modern human one in more than just shape [69<sup>\*\*</sup>].

Neanderthals had elaborate cultural adaptations that were virtually identical to AMH (until the latter developed the Upper Paleolithic toolkit after the demise of the Neanderthals), they flexibly adapted to their ecologies, buried their dead (apparently with mortuary ceremonies), built large structures, penetrated deeply into caves, and seem to have adorned themselves with drilled beads and feathers. They inhabited a climatic range (from Arctic to Mediterranean) requiring the extended ability to use culture as a buffer for ecological adaptation, and they interacted repeatedly with AMH over tens of thousands of years, exchanging genes, parasites and culture.

It will always be hard to prove the existence of prehistoric language and to infer its properties beyond the 10ky range of historical linguistics [70]. Almost no-one doubts AMH had language 60 kya just before the great diaspora, since all branches of humankind have language, but how far back should we project it? The evolution of language involved very probably a complex interplay between



exaptation, reorganization, phenotypic plasticity, adaptation and cultural evolution, touching many components and processes, requiring tens or hundreds of thousands of years and proceeding at highly variable rates [1,71–73]. The ‘externalization’ of language and its neural underpinnings, far from being trivial, is, in our view, one of the clearest clues to its long gestation. Speech itself involves over 100 muscles and complex anatomical structures that require coordination and planning at the ten-milliseconds and millimeter scales, and the evolution of specialized neural connections to the tongue, the larynx and the intercostal muscles, the extension of the arcuate fasciculus and the development of other neural circuits (e.g. [74]). A deep prehistory to language seems inevitable, especially as the new analyses of early AMH fossils in Morocco already double the time depth of our own lineage [3\*,40\*], and part of this prehistory may be traceable in the interactional infrastructure for language which includes the sustained mutual attention and the exchange of turns at production, infrastructure which is probably phylogenetically deep among the primates [75].

Language seems to be a precondition for the transmission of elaborate lithic technologies: novices were systematically taught [65] the simple *Oldowan* technology under five different conditions (reverse engineering, emulation, basic teaching, gestural teaching and spoken teaching) and found that full linguistic teaching led to the most efficient technique and good tools, followed by gestural teaching. The reasons are that efficient knapping relies on not easily observable skills, like knowledge about platform angles just under 90 degrees. If one considers all of the cultural skills needed to survive in ecologies from the Arctic to game-poor Mediterranean littorals, it is difficult to argue that Neanderthals lacked complex linguistic codes, capable of communicating about spatial locations, hunting and gathering, fauna and flora, social relations, technologies, and so on. This would imply a large lexicon, and propositional encoding. Granting Neanderthals advanced language capacities seems to us inevitable.

## Conflict of interest statement

Nothing declared.

## Acknowledgements

We wish to thank Ludy Cilissen for help with drawing the figure, Cedric Boeckx and Didier Demolin for discussion, and William Marslen-Wilson for comments on an earlier draft. DD was funded by the NWO VIDI Grant No. 276-70-022 and an EURIAS Grant 2017–2018.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Dediu D, Levinson SC: **On the antiquity of language: the reinterpretation of Neanderthal linguistic capacities and its consequences.** *Front Psychol* 2013, **4**:397.

2. Berwick RC, Chomsky N: **Why only us: recent questions and answers.** *J Neurolinguist* 2017, **43(Part B)**:166–177.

3. Hublin J-J, Ben-Ncer A, Bailey SE, Freidline SE, Neubauer S, Skinner MM, Bergmann I, Le Cabec A, Benazzi S, Harvati K *et al.*: **New fossils from Jebel Irhoud, Morocco and the pan-African origin of Homo sapiens.** *Nature* 2017, **546**:289–292.

It extends both the range (to the whole of Africa) and time depth (c. 300 kya) of modern human evolution; moreover, these AMH show an intriguing mosaic of features with a fully modern face but an archaic, non-globular cranium.

4. Meyer M, Arsuaga J-L, de Filippo C, Nagel S, Aximu-Petri A, Nickel B, Martínez I, Gracia A, de Castro JMB, Carbonell E *et al.*: **Nuclear DNA sequences from the Middle Pleistocene Sima de los Huesos hominins.** *Nature* 2016, **531**:504–507.

5. Vernot B, Tucci S, Kelso J, Schraiber JG, Wolf AB, Gittelman RM, Dannemann M, Grote S, McCoy RC, Norton H *et al.*: **Excavating Neanderthal and Denisovan DNA from the genomes of Melanesian individuals.** *Science* 2016 <http://dx.doi.org/10.1126/science.aad9416>.

This paper uses innovative methods to extract information about extinct archaic humans from the genomes of present-day humans in a ‘fossil-free manner’, including on interbreeding and the selective pressures affecting ancient DNA in the modern human genomic context.

6. Liu W, Martínón-Torres M, Cai Y, Xing S, Tong H, Pei S, Sier MJ, Wu X, Edwards RL, Cheng H *et al.*: **The earliest unequivocally modern humans in southern China.** *Nature* 2015, **526**:696–699.

7. Kuhlwlrm M, Gronau I, Hubisz MJ, de Filippo C, Prado-Martinez J, Kircher M, Fu Q, Burbano HA, Lalueza-Fox C, de la Rasilla M *et al.*: **Ancient gene flow from early modern humans into Eastern Neanderthals.** *Nature* 2016, **530**:429–433.

8. Trinkaus E, Moldovan O, Milota S, Bilgâr A, Sarcina L, Athreya S, Bailey SE, Rodrigo R, Mircea G, Higham T *et al.*: **An early modern human from the Peștera cu Oase, Romania.** *Proc Natl Acad Sci U S A* 2003, **100**:11231–11236.

9. Fu Q, Hajdinjak M, Moldovan OT, Constantin S, Mallick S, Skoglund P, Patterson N, Rohland N, Lazaridis I, Nickel B *et al.*: **An early modern human from Romania with a recent Neanderthal ancestor.** *Nature* 2015, **524**:216–219.

10. Posth C, Wißing C, Kitagawa K, Paganí L, van Holstein L, Racimo F, Wehrberger K, Conard NJ, Kind CJ, Bocherens H *et al.*: **Deeply divergent archaic mitochondrial genome provides lower time boundary for African gene flow into Neanderthals.** *Nat Commun* 2017, **8**:16046.

11. Sankararaman S, Mallick S, Dannemann M, Prüfer K, Kelso J, Pääbo S, Patterson N, Reich D: **The genomic landscape of Neanderthal ancestry in present-day humans [Internet].** *Nature* 2014. (advance online publication).

See annotation to Ref. [5\*\*].

12. Sankararaman S, Mallick S, Patterson N, Reich D: **The combined landscape of Denisovan and Neanderthal ancestry in present-day humans.** *Curr Biol* 2016, **26**:1241–1247.

See annotation to Ref. [5\*\*].

13. Weyrich LS, Duchene S, Soubrier J, Arriola L, Llamas B, Breen J, Morris AG, Alt KW, Caramelli D, Dresely V *et al.*: **Neanderthal behaviour, diet, and disease inferred from ancient DNA in dental calculus.** *Nature* 2017, **544**:357–361.

14. Pimenoff VN, Oliveira D, Mendes C, Bravo IG: **Transmission between archaic and modern human ancestors during the evolution of the oncogenic human papillomavirus 16.** *Mol Biol Evol* 2017, **34**:4–19.

15. Nielsen R, Akey JM, Jakobsson M, Pritchard JK, Tishkoff S, Willerslev E: **Tracing the peopling of the world through genomics.** *Nature* 2017, **541**:302–310.

Comprehensive and up-to-date review of human expansion around the world using genomic data and methods.

16. Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, Heinze A, Renaud G, Sudmant PH, de Filippo C *et al.*: **The complete genome sequence of a Neanderthal from the Altai Mountains.** *Nature* 2014, **505**:43–49.

17. Vernot B, Akey JM: **Resurrecting surviving Neandertal lineages from modern human genomes**. *Science* 2014 <http://dx.doi.org/10.1126/science.1245938>.  
See annotation to Ref. [5\*\*].
18. Fisher SE, Vernes SC: **Genetics and the language sciences**. *Annu Rev Linguist* 2015, **1**:289-310.
19. Devanna P, Dediu D, Vernes SC: **Language and genetics**. In *Oxford Handbook of Psycholinguistics*. Edited by Gaskell G, Rueschmeyer S-A. Oxford University Press; 2018. (in press).
20. Maricic T, Günther V, Georgiev O, Gehre S, Cürlin M, Schreiweis C, Naumann R, Burbano HA, Meyer M, Lalueza-Fox C *et al.*: **A recent evolutionary change affects a regulatory element in the human FOXP2 gene**. *Mol Biol Evol* 2013, **30**:844-852.
21. Mozzi A, Forni D, Clerici M, Pozzoli U, Mascheretti S, Guerini FR, Riva S, Bresolin N, Cagliani R, Sironi M: **The evolutionary history of genes involved in spoken and written language: beyond FOXP2**. *Sci Rep* 2016, **6**.
22. Dannemann M, Prüfer K, Kelso J: **Functional implications of Neandertal introgression in modern humans**. *Genome Biol* 2017, **18**:61.  
See annotation to Ref. [26\*\*].
23. Harris K, Nielsen R: **The genetic cost of Neandertal introgression**. *Genetics* 2016, **203**:881-891.  
It explores the selection against archaic DNA in the modern human genomic context and proposes that most of this might be due to the much smaller archaic effective population sizes that resulted in an accumulation of slightly deleterious alleles.
24. Fu Q, Posth C, Hajdinjak M, Petr M, Mallick S, Fernandes D, Furtwängler A, Haak W, Meyer M, Mittnik A *et al.*: **The genetic history of Ice Age Europe [Internet]**. *Nature* 2016. (advance online publication).
25. Jégou B, Sankararaman S, Rolland AD, Reich D, Chalmel F: **Meiotic genes are enriched in regions of reduced archaic ancestry**. *Mol Biol Evol* 2017 <http://dx.doi.org/10.1093/molbev/msx141>.  
See annotation to Ref. [26\*\*].
26. McCoy RC, Wakefield J, Akey JM: **Impacts of Neandertal-introgressed sequences on the landscape of human gene expression**. *Cell* 2017, **168** 916-927.e12.  
This paper try to understand the functional effects of introgressed archaic DNA in the modern human genomes using a large array of methods ranging from electronic healthcare records [27] to gene expression [22,26\*\*].
27. Simonti CN, Vernot B, Bastarache L, Bottinger E, Carrell DS, Chisholm RL, Crosslin DR, Hebbbring SJ, Jarvik GP, Kullo IJ *et al.*: **The phenotypic legacy of admixture between modern humans and Neanderthals**. *Science* 2016, **351**:737-741.  
See annotation to Ref. [26\*\*].
28. Juric I, Aeschbacher S, Coop G: **The strength of selection against Neandertal introgression**. *PLOS Genet* 2016, **12**: e1006340.  
See annotation to Ref. [23\*].
29. Sams AJ, Dumaine A, Nédélec Y, Yotova V, Alfieri C, Tanner JE, Messer PW, Barreiro LB: **Adaptively introgressed Neandertal haplotype at the OAS locus functionally impacts innate immune responses in humans**. *Genome Biol* 2016, **17**:246.
30. Dannemann M, Andrés AM, Kelso J: **Introgression of Neandertal- and Denisovan-like haplotypes contributes to adaptive variation in human toll-like receptors**. *Am J Hum Genet* 2016, **98**:22-33.
31. Deschamps M, Laval G, Fagny M, Itan Y, Abel L, Casanova J-L, Patin E, Quintana-Murci L: **Genomic signatures of selective pressures and introgression from archaic hominins at human innate immunity genes**. *Am J Hum Genet* 2016, **98**:5-21.
32. Gittelman RM, Schraiber JG, Vernot B, Mikacenic C, Wurfel MM, Akey JM: **Archaic hominin admixture facilitated adaptation to out-of-Africa environments**. *Curr Biol* 2016, **26**:3375-3382.
33. Quach H, Rotival M, Pothlichet J, Loh Y-HE, Dannemann M, Zidane N, Laval G, Patin E, Harmant C, Lopez M *et al.*: **Genetic adaptation and Neandertal admixture shaped the immune system of human populations**. *Cell* 2016, **167** 643-656.e17.
34. Ding Q, Hu Y, Xu S, Wang J, Jin L: **Neandertal introgression at chromosome 3p21.31 was under positive natural selection in East Asians**. *Mol Biol Evol* 2014, **31**:683-695.
35. Quach H, Quintana-Murci L: **Living in an adaptive world: genomic dissection of the genus Homo and its immune response**. *J Exp Med* 2017, **214**:877-894.  
Up-to-date, comprehensive and very readable review of the genetic bases of immunity and the relevance of admixture with archaic humans in introducing adaptive variants into the modern human genome.
36. Boeckx CA, Benítez-Burraco A: **The shape of the human language-ready brain**. *Front Psychol* 2014, **5**:282.
37. Boeckx C, Benítez-Burraco A: **Globularity and language-readiness: generating new predictions by expanding the set of genes of interest [Internet]**. *Front Psychol* 2014, **5**.
38. Benítez-Burraco A, Boeckx C: **Possible functional links among brain- and skull-related genes selected in modern humans [Internet]**. *Front Psychol* 2015, **6**.
39. Benítez-Burraco A, Theofanopoulou C, Boeckx C: **Globalization and domestication**. *Topoi* 2016 <http://dx.doi.org/10.1007/s11245-016-9399-7>.
40. Richter D, Grün R, Joannes-Boyau R, Steele TE, Amani F, Rué M, Fernandes P, Raynal J-P, Geraads D, Ben-Ncer A *et al.*: **The age of the hominin fossils from Jebel Irhoud, Morocco, and the origins of the Middle Stone Age**. *Nature* 2017, **546**:293-296.  
See annotation to Ref. [3\*].
41. Lieberman P: **Vocal tract anatomy and the neural bases of talking**. *J Phon* 2012, **40**:608-622.
42. Lieberman P: **The evolution of language and thought**. *J Anthropol Sci* 2016, **94**:127-146.
43. Gokhman D, Lavi E, Prüfer K, Fraga MF, Riancho JA, Kelso J, Pääbo S, Meshorer E, Carmel L: **Reconstructing the DNA methylation maps of the Neandertal and the Denisovan**. *Science* 2014, **344**:523-527.
44. Roebroeks W, Soressi M: **Neandertals revised**. *Proc Natl Acad Sci U S A* 2016, **113**:6372-6379.
45. Soressi M, McPherron SP, Lenoir M, Dogandzic T, Goldberg P, Jacobs Z, Maigrot Y, Martisius NL, Miller CE, Rendu W *et al.*: **Neandertals made the first specialized bone tools in Europe**. *Proc Natl Acad Sci* 2013, **110**:14186-14190.
46. Wales N: **Modeling Neandertal clothing using ethnographic analogues**. *J Hum Evol* 2012, **63**:781-795.
47. Sørensen B: **Energy use by Eem Neanderthals**. *J Archaeol Sci* 2009, **36**:2201-2205.
48. Shipley GP, Kindscher K: **Evidence for the paleoethnobotany of the Neandertal: a review of the literature [Internet]**. *Scientifica (Cairo)* 2016.
49. El Zaatari S, Grine FE, Ungar PS, Hublin J-J: **Neandertal versus modern human dietary responses to climatic fluctuations**. *PLOS ONE* 2016, **11**.
50. Estalrich A, El Zaatari S, Rosas A: **Dietary reconstruction of the El Sidrón Neandertal familial group (Spain) in the context of other Neandertal and modern hunter-gatherer groups. A molar microwear texture analysis**. *J Hum Evol* 2017, **104**:13-22.
51. Krueger KL, Ungar PS, Guatelli-Steinberg D, Hublin J-J, Pérez-Pérez A, Trinkaus E, Willman JC: **Anterior dental microwear textures show habitat-driven variability in Neandertal behavior**. *J Hum Evol* 2017, **105**:13-23.
52. Hardy K, Buckley S, Collins MJ, Estalrich A, Brothwell D, Copeland L, García-Tabernero A, García-Vargas S, Rasilla Mde la, Lalueza-Fox C *et al.*: **Neandertal medics? Evidence for food, cooking, and medicinal plants entrapped in dental calculus**. *Naturwissenschaften* 2012, **99**:617-626.
53. Frayer DW, Gatti J, Monge J, Radović D: **Prehistoric dentistry? P4 rotation, partial M3 impaction, toothpick grooves and other**

- signs of manipulation in Krapina Dental Person 20.** *Bull Int Assoc Paleodontology* 2017, **11**:1-10.
54. Welker F, Hajdinjak M, Talamo S, Jaouen K, Dannemann M, David F, Julien M, Meyer M, Kelso J, Barnes I *et al.*: **Palaeoproteomic evidence identifies archaic hominins associated with the Châtelperronian at the Grotte du Renne.** *Proc Natl Acad Sci U S A* 2016, **113**:11162-11167.
- Association of the transitional Châtelperronian at the Grotte du Renne with the Neanderthals using proteomic and genetic methods.
55. Muller A, Clarkson C: **Identifying major transitions in the evolution of lithic cutting edge production rates.** *PLOS ONE* 2016, **11**.
  56. Rougier H, Crevecoeur I, Beauval C, Posth C, Flas D, Wißing C, Furtwängler A, Germonpré M, Gómez-Olivencia A, Semal P *et al.*: **Neanderthal cannibalism and Neanderthal bones used as tools in Northern Europe.** *Sci Rep* 2016, **6**.
  57. Villa P, Soriano S, Grün R, Marra F, Nomade S, Pereira A, Boschian G, Pollarolo L, Fang F, Bahain J-J: **The Acheulian and Early Middle Paleolithic in Latium (Italy): stability and innovation.** *PLOS ONE* 2016, **11**.
  58. Heyes PJ, Anastasakis K, de Jong W, van Hoesel A, Roebroeks W, Soressi M: **Selection and use of manganese dioxide by Neanderthals.** *Sci Rep* 2016, **6**.
  59. Mellars P: **Neanderthal symbolism and ornament manufacture: the bursting of a bubble?** *PNAS* 2010, **107**:20147-20148.
  60. Finlayson C, Brown K, Blasco R, Rosell J, Negro JJ, Bortolotti GR, Finlayson G, Sánchez Marco A, Giles Pacheco F, Rodríguez Vidal J *et al.*: **Birds of a feather: Neanderthal exploitation of raptors and corvids.** *PLoS ONE* 2012, **7**:e45927.
  61. Majkić A, Evans S, Stepanchuk V, Tsvetikh A, d'Errico F: **A decorated raven bone from the Zaskalnaya VI (Kolosovskaya) Neanderthal site, Crimea.** *PLOS ONE* 2017, **12**:e0173435.
  62. Jaubert J, Verheyden S, Genty D, Soulier M, Cheng H, Blamart D, Burlet C, Camus H, Delaby S, Deldicque D *et al.*: **Early Neanderthal constructions deep in Bruniquel Cave in southwestern France.** *Nature* 2016, **534**:111-114.
  63. Austin P, Bresnan J: **Non-configurationality in Australian aboriginal languages.** *Nat Lang Linguist Theory* 1996, **14**:215-268.
  64. Stout D: **Stone toolmaking and the evolution of human culture and cognition.** *Philos Trans R Soc Lond B Biol Sci* 2011, **366**:1050-1059.
  65. Morgan TJH, Uomini NT, Rendell LE, Chouinard-Thuly L, Street SE, Lewis HM, Cross CP, Evans C, Kearney R, de la Torre I *et al.*: **Experimental evidence for the co-evolution of hominin tool-making teaching and language.** *Nat Commun* 2015, **6**:6029.
  66. Wynn T, Overmann KA, Coolidge FL: **The false dichotomy: a refutation of the Neanderthal indistinguishability claim.** *J Anthropol Sci* 2016, **94**:1-22.
  67. Beals ME, Frayer DW, Radović J, Hill CA: **Cochlear labyrinth volume in Krapina Neandertals.** *J Hum Evol* 2016, **90**:176-182.
  68. Stoessel A, David R, Gunz P, Schmidt T, Spoor F, Hublin J-J: **Morphology and function of Neanderthal and modern human ear ossicles.** *Proc Natl Acad Sci U S A* 2016, **113**:11489-11494.
- This paper analyses in detail the largest available sample of Neanderthal ear ossicles and concludes that despite morphological differences from AMH, they were functionally very similar, pointing to selective pressures to maintain 'consistent' vocal communication in the two lineages.
69. D'Anastasio R, Wroe S, Tuniz C, Mancini L, Cesana DT, Dreossi D, Ravichandiran M, Attard M, Parr WCH, Agur A *et al.*: **Micro-biomechanics of the Kebara 2 hyoid and its implications for speech in Neanderthals.** *PLoS ONE* 2013, **8**:e82261.
- Modelling showing that the Neanderthal hyoid was not only morphologically but also biomechanically very similar to the modern human one.
70. Renfrew C, McMahon A, Trask L (Eds): *Time Depth in Historical Linguistics*. UK: McDonald Institute; 2000.
  71. Fitch WT: **Evolutionary developmental biology and human language evolution: constraints on adaptation.** *Evol Biol* 2012, **39**:613-637.
  72. Laland KN, Uller T, Feldman MW, Sterelny K, Müller GB, Moczek A, Jablonka E, Odling-Smee J: **The extended evolutionary synthesis: its structure, assumptions and predictions.** *Proc R Soc B* 2015, **282**:20151019.
  73. Dediu D, Cysouw M, Levinson SC, Baronchelli A, Christiansen MH, Croft W, Evans N, Garrod S, Gray R, Kandler A *et al.*: **Cultural evolution of language.** In *Cultural Evolution: Society, Technology, Language, and Religion*. Edited by Richerson PJ, Christiansen MH. MIT Press; 2013:303-332.
  74. Simonyan K, Horwitz B: **Laryngeal motor cortex and control of speech in humans.** *Neuroscientist* 2011, **17**:197-208.
  75. Levinson SC: **Turn-taking in human communication – origins and implications for language processing.** *Trends Cogn Sci* 2016, **20**:6-14.